TIMED Solar EUV Experiment:

Phase E Annual Report for 2002

SEE Science Team

LASP/CU: Tom Woods (PI), Frank Eparvier, Don Woodraska, Gary Rottman

HAO/NCAR: Stan Solomon, Ray Roble, Giuliana de Toma, Dick White

NRL: Judith Lean

SpaceWx: Kent Tobiska

GI/UAF: Scott Bailey

Report Outline

- ◆ SEE Science Overview

- ◆ SEE Instrument Overview and Status

- ◆ Summary of SEE Data Products

- ◆ Summary of SEE Results

- ◆ Summary of SEE Related Talks and Papers

- ◆ Future Plans for SEE Team

SEE Science Plans

Solar UV Irradiance Measurements

Validations
Internal Calibrations,
Underflight Calibrations
SOHO, SNOE,
UARS, SORCE

Eparvier, Woods, Bailey, Rottman

Solar UV Variability
Function of wavelength
Over time scales of minutes to years

All

Study Earth's Response
Photoelectron analysis with FAST data
and using the *glow* model
Atmospheric response studies using
HAO's **TIM-GCM**

Solomon, Roble, Bailey, Eparvier

Modeling Solar Variation
Study variations related to active region
evolution derived from solar images
Improve the **NRLEUV, SOLAR2000,**
and **SunRise** solar irradiance models

Lean, Tobiska, White, Fox, Woods

TIMED Solar EUV Experiment

MU   Radiator

SSPP
Actuator

XPS

SAS   EGS

Measures the solar vacuum
ultraviolet (VUV) irradiance
Range:
 0.1-194 nm
Resolution:
 0.4 nm EGS (27-194 nm)
 5-10 nm XPS (0.1-34 nm)
Frequency:
 10-sec integrations, but only
 for 3 min per orbit (96 min)

MU = Microprocessor Unit
SSPP = SEE Solar Pointing Platform
SAS = Solar Aspect Sensor (2)

EGS = EUV Grating Spectrograph
Rowland-circle grating spectrograph with
64x1024 CODACON (MCP-based) detector

Uses 2 slits to provide redundant measurements

XPS = XUV Photometer System
Set of 12 Si photodiodes - 8 for XUV, 1 for Ly-α,
and 3 for window calibrations

Includes 3 redundant photodiodes

LASP

Status of SEE Instrument

- **EUV Grating Spectrograph (EGS)**
  - Vacuum door anomaly on 2002/005, solved on 2002/011
  - Fully functional instrument
  - Some degradation, mostly at the bright lines on the CODACON (MCP-based) detector, but is tracked well with on-board redundant channel and flat-field detector lamp weekly experiments
- **XUV Photometer System (XPS)**
  - Fully functional until 2002/205 when there was a filter wheel anomaly (filter wheel stuck in position 6)
  - 3 channels continue to make good solar measurements
  - SORCE, with identical XPS, launches in Jan. 2003
  - No detectable degradation, except for slight degradation for the Lyman-α channel, as determined by redundant channels used up to 2002/205
- **Microprocessor Unit (MU)**
  - Fully functional
- **SEE Solar Pointing Platform (SSPP)**
  - Fully functional

LASP

TIMED Mission Began at Solar Maximum

- TIMED was just in time for solar maximum
- Solar cycle 23 is now dropping towards solar minimum
- SEE's daily observations began on Jan. 22, 2002

Summary of SEE Observations

- Results are for the period of Jan. 22 - Dec. 11, 2002
- Observations in 2002
  - 4,798 orbits (96%) with solar observations
  - 322 days (99.4%) with EGS observations
  - 318 days (98.1%) with XPS observations
  - 184 calibration (redundant channel) solar observations
  - 149 EGS detector flat-field calibrations (Hg lamp)
  - Special underflight calibration rocket
    - NASA 36.192 launched on Feb. 8, 2002, complete success
- Data Gaps in 2002
  - January 1-21: still in instrument commissioning phase
  - March 2: S/C safehold (yaw around error)
  - March 4: gap for EGS only due to improper slit command (ground S/W error)
  - July 25-29: gap for XPS only due to its filter wheel anomaly

Status of SEE Data Processing

- ♦ SEE data products are being generated daily
  - Version 5 is current version
- ♦ SEE XPS and EGS Level 2 data products are available now
  - Daily average, instrument resolution, atmospheric absorption corrected, degradation corrected, normalized to 1 AU
  - NetCDF format: use IDL read_netcdf.pro to read SEE data products
  - Quick-look: use IDL plotxps_ts.pro or plotegs_ts.pro to plot time series
- ♦ SEE Level 3 data products will be available in Version 6 release
  - SEE Level 3 is the solar irradiance in 1 nm bins on 0.5 nm centers from 0-195 nm along with a list of the irradiances of the brighter emission lines
  - EGS data is used above 27 nm and XPS data with a solar model is used below 27 nm
  - These Version 6 data will be a more fully *validated* data set. Version 5 data have calibration issues related to broadband sensitivity for XPS and order sorting for EGS. Release of SEE Version 6 is expected in early 2003.

Summary of SEE Data Products

- ♦ Summary is for all versions
- ♦ SEE Low Level (raw) Data Products (file per day)
  - Level 0A (TP) files: 389
- ♦ XPS Data Products (file per day)
  - Level 0B files: 682
  - Level 1 files: 702
  - Level 2 files (access to public): 813
- ♦ EGS Data Products (file per day)
  - Level 0B files: 1110
  - Level 1 files: 711
  - Level 2 files (access to public): 1433
- ♦ SEE Level 3 (1 nm merged) Data Products (file per day)
  - Level 3 files: 717
  - Will be made public for SEE Version 6 data products

SEE Validation Results

SNOE/TIMED 2-7 nm Comparison
TIMED is ~30% lower than SNOE

SNOE SXP
TIMED SEE
Rocket

UARS SOLSTICE / EGS

Average difference < 1%

- ◆ Good agreement between XPS and SNOE below 17 nm
- ◆ Excellent agreement between EGS and UARS above 119 nm
- ◆ Issues to be resolved in Version 6 data are:
  - • XPS 17-21 nm issue (0-7 nm contamination for the Al/Nb photometer)
  - • EGS 26-115 nm issue (higher grating order correction for rocket EGS)

Reminder of SEE Science Objectives

1. Accurately and precisely determine the time-dependent solar vacuum ultraviolet (VUV: below 200 nm) spectral irradiance

2. Study the solar-terrestrial relationships utilizing atmospheric models, primarily the TIME-GCM at HAO/NCAR

3. Study solar VUV variability (27-day rotations, solar cycle changes) and its sources

4. Improve proxy models of the solar VUV irradiance

5. Determine the thermospheric neutral densities (O_2, N_2 and O) from solar occultations

Summary of SEE Results

- ♦ **Objective 1: solar VUV spectral irradiance measurements**
  - Daily measurements since Jan. 22, 2002
  - On-going validation effort to verify 10-20% accuracy and 2-4% precision
- ♦ **Objective 2: model solar response in Earth's atmosphere**
  - Use of *glow* model with SEE solar data and FAST photoelectron data
- ♦ **Objective 3: solar variability**
  - Several solar rotations (11) observed during solar maximum conditions
  - Many flares (> 30) observed by SEE during TIMED mission
    - New, unexpected results from SEE !
- ♦ **Objective 4: solar irradiance modeling**
  - NRLEUV model improvements: extended below 5 nm and added flare DEM
  - SOLAR2000 model improvements: extended above 122 nm to 200 nm
- ♦ **Objective 5: densities from solar occultation**
 - Several occultation observations obtained, but only preliminary analysis to date

---

Example Solar VUV (0-200 nm) Spectrum

Example Solar Variations from XPS

XPS has observed over 30 flares, including 3 large X-class flares.
XUV variations are a factor of about 3 during 2002.
Lyman-α variations are less, as expected, at about 40%.

There are now only 3 solar measurements from XPS since the XPS filter wheel anomaly on day 2002/205 (July 24).

The XPS on the SORCE satellite will start a full set of XUV measurements in Feb. 2003.

Shown are the XPS Level 2 data (daily averaged)

Example Solar Variations from EGS

Special Study is Large Flare on April 21, 2002

- X-ray increased by factor of 8, **peak appears to be near 2 nm**
- Longer wavelengths enhanced much less (10-40%)

The SEE measurements are key for the study of the solar photon energy input during the April storms.

- X1.5 class flare at ~2 UT
- C8 class flare at ~18 UT

XPS
FAST e⁻
NRLEUV
X-rays and coronal lines vary the most
EGS

Flare Ratio — LOG Scale

Wavelength (nm): 0   50   100   150   200

Results of SEE solar irradiance measurements and FAST photoelectron measurements are given in Woods et al. [GRL, submitted, 2002].

Relative Irradiance

FAST   X
            M
GOES 0.1-0.8 nm
            C
SEE 0.1-7 nm

Time (hour): 0   5   10   15   20

LASP   **TIMED SEE**   *SEE Annual Report Jan. 2003 - 16*

Modeling Photoelectron Data

- Obtained FAST photoelectron data (shown as the square symbols) at nearly the same time as the SEE measurements during the April 21, 2002 X-class flare event. The dashed line is the photoelectron background signal level that has already been removed.
- The *glow* model (results shown as the solid lines) uses the SEE solar irradiances as input
- The *glow* model result for the pre-flare condition is in good agreement with the photoelectron data
- The glow model result for near the flare peak is higher than the data, partially because the SEE solar irradiance is closer to the flare peak than the photoelectron data

Flux (e⁻ cm⁻² s⁻¹ sr⁻¹ eV⁻¹)

Energy (eV): 10   100   1000

□ Pre-flare FAST photoelectron flux
□ Flare FAST photoelectron flux
── Pre-flare *glow* model result
── Flare *glow* model result

LASP   **TIMED SEE**   *SEE Annual Report Jan. 2003 - 17*

New Measurements of the Solar Rotation Variability

Solar Cycle 23

SEE has observed 10 solar rotations so far.

Choose a particularly strong
solar rotation for comparison
study: Day 220 to Day 232
(Aug. 8 to Aug. 20, 2002)

TIMED Mission

Rotational
Variability
Study

Day
220

Day
232

Day of Year 2002

LASP

TIMED SEE

SEE Annual Report Jan. 2003 - 18

Example Spectrum of Solar Rotation Variability

27-Day Variability

XPS 1 nm bins

EGS 1 nm bins

EGS lines

Reference
(Woods & Rottman, 2002)

Reference is based mostly
on UARS in the FUV
and on AE-E in the EUV

50-90%
variability
in XUV

Ratio of Days 232/220

10-40% variability
in EUV lines and continua

1-15% variability in FUV

Wavelength (nm)

LASP

TIMED SEE

SEE Annual Report Jan. 2003 - 19

Classification of Solar Emissions

TIMED SEE (preliminary results)   UARS SOLSTICE

- ◆ **Classify emission type by:**
  - • Photosphere
  - • Chromosphere
  - • Transition Region
  - • Corona

3-Parameter Model with Mg II C/W Index

$$E = E_{min}\left(1 + C_S \frac{Mg - Mg_{54}}{Mg_{min}} + C_L \frac{Mg_{54} - Mg_{min}}{Mg_{min}}\right)$$

E = Irradiance Mg = MgII C/W ratio

C_S = Contrast ratio for Short – term (< 54 days)

C_L = Contrast ratio for Long – term (> 54 days)

Mg_{54} = Mg smoothed 54 days (prior to current day)

LASP   **TIMED SEE**   *SEE Annual Report Jan. 2003 - 20*

SEE Related Talks in 2002

- ◆ TIMED Science Meeting: Feb 2002, 3 talks
- ◆ AGU Spring Meeting: May 2002, 4 talks
- ◆ SORCE Science Meeting: July 2002, 1 talk
- ◆ April Storm Workshop: Aug 2002, 1 talk
- ◆ COSPAR Meeting: Oct 2002, 1 talk
- ◆ NOAA Solar EUV Workshop: Oct 2002, 1 talk
- ◆ AGU Fall Meeting: Dec 2002, 4 talks
- ◆ Public Seminars: LASP/CU, NOAA, NCAR, Michigan Tech. Univ.

LASP   **TIMED SEE**   *SEE Annual Report Jan. 2003 - 21*

SEE Related Papers in 2002

- Woods, T. N., S. M. Bailey, W. K. Peterson, H. P. Warren, S. C. Solomon, F. G. Eparvier, H. Garcia, C. W. Carlson, and J. P. McFadden, Solar extreme ultraviolet variability of the X-class flare on April 21, 2002 and the terrestrial photoelectron response, GRL, submitted, 2002.

- Eparvier, F. G., T. N. Woods, D.L. Woodraska, S.M. Bailey, and S.C. Solomon, Spectral irradiance measurements from the TIMED Solar EUV Experiment, *Advances in Space Research*, submitted, 2002.

- Woods, T. N., F. G. Eparvier, S. C. Solomon, D. L. Woodraska, and S. M. Bailey, Early results from the TIMED Solar EUV Experiment, *Proceedings of 4th TIGER Symposium*, virtual journal at http://www.ipm.fraunhofer.de/english/meetings/workshops/tiger/, June 2002.

- Tobiska, W.K., New developments in solar irradiance proxies for operational space weather, *Proceedings of 4th TIGER Symposium*, virtual journal at http://www.ipm.fraunhofer.de/english/meetings/workshops/tiger/, June 2002.

- Woods, T., L. W. Acton, S. Bailey, F. Eparvier, H. Garcia, D. Judge, J. Lean, D. McMullin, G. Schmidtke, S. C. Solomon, W. K. Tobiska, and H. P. Warren, Solar extreme ultraviolet and x-ray irradiance variations, in *Solar Variability and Its Effect on Earth's Atmospheric and Climate System*, eds. J. Pap, C. Fröhlich, H. Hudson, J. Kuhn, J. McCormack, G. North, W. Sprig, and S. T. Wu, Geophys. Monograph Series, Wash. DC, in press, 2002.

- Thuillier, G., T. N. Woods, L. E. Floyd, R. Cebula, M. Hersé, and D. Labs, Reference solar spectra during solar cycle 22, in *Solar Variability and Its Effect on Earth's Atmospheric and Climate System*, eds. J. Pap, C. Fröhlich, H. Hudson, J. Kuhn, J. McCormack, G. North, W. Sprig, and S. T. Wu, Geophys. Monograph Series, Wash. DC, in press, 2002.

- Woods, T. and G. Rottman, Future solar irradiance observations from the NASA TIMED and SORCE satellites, in *Radiometric Inter-calibration of SOHO*, eds.M. C. E. Huber, A. Pauluhn, and R. von Steiger, Bern, Switz., in press, 2002.

- McMullin, D., T. Woods, I. E. Dammasch, D. Judge, P. Lemaire, J. S. Newmark, W. Thompson, W. K. Tobiska, and K. Wilhelm, Irradiance working group report for the SOHO intercalibration workshop, in *Radiometric Inter-calibration of SOHO*, eds.M. C. E. Huber, A. Pauluhn, and R. von Steiger, Bern, Switz., in press, 2002.

- Woods, T. N. and G. J. Rottman, Solar ultraviolet variability over time periods of aeronomic interest, in *Comparative Aeronomy in the Solar System*, eds. M. Mendillo, A. Nagy, and J. Hunter Waite, Jr., Geophys. Monograph Series, Wash. DC, pp. 221-234, 2002.

SEE Future Plans

- ## Complete additional validations
  - Validated SEE Version 6 data should be available in early 2003

- ## Detailed solar variability studies
  - More complete analysis of the 11+ solar rotation and 30+ flare observations
  - Improve models of the solar UV irradiance

- ## Detailed modeling of Earth's response to solar irradiance changes
  - Composition, dynamics, temperature using TIME-GCM
  - Photoelectrons using *glow* model

- ## Occultation data analysis